

## Factoring

- $a^2 - b^2 = (a - b)(a + b)$
- $a^2 + b^2$  is prime
- $a^2 + 2ab + b^2 = (a + b)^2$
- $a^2 - 2ab + b^2 = (a - b)^2$
- $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
- $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

## Analytic Geometry

- slope:  $m = \frac{y_2 - y_1}{x_2 - x_1}$
- equation of a line:  $y - y_1 = m(x - x_1)$
- distance:  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

## Exponent Rules

- $a^{x+y} = a^x a^y$
- $(ab)^x = a^x b^x$
- $(a^x)^y = a^{xy}$
- $a^0 = 1$  if  $a \neq 0$
- $a^{-x} = \frac{1}{a^x}$  if  $a \neq 0$
- $a^{x-y} = \frac{a^x}{a^y}$  if  $a \neq 0$

## Logarithm Rules

- $\log_b x = y \iff x = b^y$
- $b^{\log_b x} = x$
- $\log_b b^x = x$
- $\log_b 1 = 0$
- $\log_b b = 1$
- $\log_b xy = \log_b x + \log_b y$
- $\log_b \frac{x}{y} = \log_b x - \log_b y$
- $\log_b x^y = y \log_b x$
- $\log_b x = \frac{\log_a x}{\log_a b}$

## Arithmetic Series

- $a_k = a + (k - 1)d$
- $S_n = \sum_{k=1}^n [a + (k - 1)d] = \frac{n}{2} [2a + (n - 1)d]$
- $S_n = \sum_{k=1}^n [a + (k - 1)d] = n \left( \frac{a + a_n}{2} \right)$

## Geometric Series

- $a_n = ar^{n-1}$
- $S_n = \sum_{k=0}^{n-1} ar^k = a \left[ \frac{1 - r^n}{1 - r} \right]$  if  $r \neq 1$
- $S = \sum_{k=0}^{\infty} ar^k = \frac{a}{1 - r}$  if  $|r| < 1$

## Trigonometry

- $\cos A = \frac{\text{adjacent}}{\text{hypotenuse}}$    ◦  $\sin A = \frac{\text{opposite}}{\text{hypotenuse}}$
- $\tan A = \frac{\text{opposite}}{\text{adjacent}}$

	0°	30°	45°	60°	90°
	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	undefined

## Pythagorean Identities

- $\cos^2 A + \sin^2 A = 1$
- $1 + \tan^2 A = \sec^2 A$
- $1 + \cot^2 A = \csc^2 A$

## Ratio Identities

- $\tan A = \frac{\sin A}{\cos A}$    ◦  $\cot A = \frac{\cos A}{\sin A}$

## Reciprocal Identities

$$\circ \sec A = \frac{1}{\cos A} \quad \circ \csc A = \frac{1}{\sin A} \quad \circ \cot A = \frac{1}{\tan A}$$

---

## Sum and Difference Identities

$$\begin{aligned} \circ \cos(A \pm B) &= \cos A \cos B \mp \sin A \sin B \\ \circ \sin(A \pm B) &= \sin A \cos B \pm \cos A \sin B \\ \circ \tan(A \pm B) &= \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \end{aligned}$$

---

## Double Angle Identities

$$\begin{aligned} \circ \cos 2A &= \cos^2 A - \sin^2 A \\ \circ \cos 2A &= 2 \cos^2 A - 1 \\ \circ \cos 2A &= 1 - 2 \sin^2 A \\ \circ \sin 2A &= 2 \cos A \sin A \\ \circ \tan 2A &= \frac{2 \tan A}{1 - \tan^2 A} \end{aligned}$$

---

## Half Angle Identities

$$\begin{aligned} \circ \cos \frac{A}{2} &= \pm \sqrt{\frac{1 + \cos A}{2}} \\ \circ \sin \frac{A}{2} &= \pm \sqrt{\frac{1 - \cos A}{2}} \\ \circ \tan \frac{A}{2} &= \frac{1 - \cos A}{\sin A} = \frac{\sin A}{1 + \cos A} \end{aligned}$$

---

## Triple Angle Identities

$$\begin{aligned} \circ \cos 3A &= 4 \cos^3 A - 3 \cos A \\ \circ \sin 3A &= 3 \sin A - 4 \sin^3 A \end{aligned}$$

---

## Power Reduction Identities

$$\begin{aligned} \circ \cos^2 A &= \frac{1 + \cos 2A}{2} \\ \circ \sin^2 A &= \frac{1 - \cos 2A}{2} \\ \circ \tan^2 A &= \frac{1 - \cos 2A}{1 + \cos 2A} \\ \circ \cos^3 A &= \frac{3 \cos A + \cos 3A}{4} \\ \circ \sin^3 A &= \frac{3 \sin A - \sin 3A}{4} \end{aligned}$$

---

## Sum-to-Product Identities

$$\begin{aligned} \circ \sin A + \sin B &= 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \\ \circ \sin A - \sin B &= 2 \cos \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right) \\ \circ \cos A + \cos B &= 2 \cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \\ \circ \cos A - \cos B &= -2 \sin \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right) \end{aligned}$$

---

## Product-to-Sum Identities

$$\begin{aligned} \circ \sin A \cos B &= \frac{1}{2} [\sin(A+B) + \sin(A-B)] \\ \circ \cos A \cos B &= \frac{1}{2} [\cos(A+B) + \cos(A-B)] \\ \circ \sin A \sin B &= \frac{1}{2} [\cos(A-B) - \cos(A+B)] \end{aligned}$$

---

## Sums of Sines and Cosines

$$\begin{aligned} \circ A \cos x + B \sin x &= \sqrt{A^2 + B^2} \sin(x + \phi) \text{ where} \\ &\cos \phi = \frac{B}{\sqrt{A^2 + B^2}} \text{ and } \sin \phi = \frac{A}{\sqrt{A^2 + B^2}} \\ \circ A \cos x + B \sin x &= \sqrt{A^2 + B^2} \cos(x - \phi) \text{ where} \\ &\cos \phi = \frac{A}{\sqrt{A^2 + B^2}} \text{ and } \sin \phi = \frac{B}{\sqrt{A^2 + B^2}} \end{aligned}$$

---

## Laws of Sines and Cosines

$$\begin{aligned} \circ c^2 &= a^2 + b^2 - 2ab \cos C \\ \circ \frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \end{aligned}$$

---

## Area of a Triangle

For a triangle with sides  $a$ ,  $b$ ,  $c$  and angles  $\angle A$ ,  $\angle B$ , and  $\angle C$ ,

$$\begin{aligned} \circ \text{Area} &= \sqrt{s(s-a)(s-b)(s-c)} \text{ where} \\ &s = \frac{a+b+c}{2} \\ \circ \text{Area} &= \frac{1}{2} ab \sin C \\ \circ \text{Area} &= \frac{c^2 \sin A \sin B}{2 \sin C} \end{aligned}$$

---

## Circular Section

$$\begin{aligned} \circ \text{Arc length: } &s = r\theta \\ \circ \text{Area: } &A = \frac{1}{2} r^2 \theta \end{aligned}$$

---